

# Helical modulation of the electrostatic potential due to magnetic islands in toroidal plasma confinement devices

**G. Ciaccio<sup>1</sup>, O. Schmitz<sup>2</sup>, S. S. Abdullaev<sup>3</sup>, T. E. Evans<sup>4</sup>,  
H. Frerichs<sup>2</sup>, G. Spizzo<sup>1</sup>, and R. B. White<sup>4</sup>**

<sup>1</sup> Consorzio RFX (CNR, ENEA, INFN, Università di Padova, Acciaierie Venete SpA), Corso Stati Uniti 4 - 35127 Padova (Italy)

<sup>2</sup> Department of Engineering Physics, University of Wisconsin - Madison, 1500 Engineering Drive, Madison, WI 53706

<sup>3</sup> Institut für Energieforschung-Plasmaphysik, Association EURATOM-FZJ, Jülich, Germany

<sup>4</sup> General Atomics, San Diego, California, USA

<sup>5</sup> Plasma Physics Laboratory, Princeton University, P.O.Box 451, Princeton, New Jersey 08543

E-mail: [giovanni.ciaccio@igi.cnr.it](mailto:giovanni.ciaccio@igi.cnr.it)

## **Abstract.**

The electrostatic response of a tokamak edge plasma to magnetic island with 4/1 poloidal/toroidal mode numbers is analyzed in direct comparison of measurements with the Hamiltonian guiding center code ORBIT. We find a strong correlation between the magnetic field topology in ion and electron velocity space and the poloidal modulation of the plasma potential measured. The ion and electron drifts yield a predominantly electron driven radial diffusion when approaching the island X-point while ion diffusivities are generally an order of magnitude smaller. This results in build up of a strong radial electric field structure pointing outward from the island O-point. An excellent agreement between measured and modeled plasma potential has been found. This shows that, in a tokamak edge plasma, a magnetic island can act as convective cell as pointed out by previous results. Here, we show for the first time that the particular drifts of electrons and ions in a 3D magnetic topology account for these effects. An analytical model for the radial particle diffusion is derived and it is shown that both, an ion and an electron diffusion dominated transport regime can exist which are known as ion and electron root solutions in stellarators. This finding and comparison to reversed field pinches shows that the role of magnetic islands as convective cells and hence as major radial particle transport drivers is generic to 3D plasma boundary layers of toroidal magnetic confinement devices.

PACS numbers: 52.20.Dq, 52.65.Cc

Plasma flows along magnetic field lines under conditions of spontaneous self-organization are a generic question in space and terrestrial plasma physics [1, 2]. In magnetically confined high temperature plasmas explored for future fusion energy production, such directed plasma flows are responsible for transport in the plasma edge. By this, the magnetic field topology in the plasma edge and the resulting transport characterizes the interface of the plasma to the surrounding neutral gas. One form of such self-organized 3D magnetic structures are magnetic islands. Field lines can be easily perturbed in a resonant way by magnetic field perturbations with the same mode structure as the rational surface of these field lines. In tokamaks, this form of resonant magnetic perturbation (RMP) is used to control plasma edge transport and stability [3, 4]. In the edge of all fusion devices, it has been observed that kinetic properties of the plasma, such as electron density and temperature [5, 6], electron pressure [7], connection length [8], in presence of 3D fields and magnetic chaos in the edge, show macroscopic modulations coherently with the symmetry of the dominant magnetic island. Moreover, it has been shown that magnetic islands influence the sign of the plasma flow,  $v$ , and the related radial electric field,  $E^r$  [7, 9, 10, 11].

The actual relation of these magnetic structures to plasma confinement and transport is an important question for fusion plasma research in particular as it has been demonstrated that magnetic islands in the plasma edge have a profound impact on plasma performance and plasma stability. At the RFP RFX-mod, for instance, a direct connection between a convective cell pattern and the empirical density limit (Greenwald limit) has been established [12, 13]. Island formation with impact on the particle confinement is also discussed as a key mechanism for stabilization of high confinement edge plasmas in tokamaks [14]. Finally, low-order rational surfaces in the periphery of stellarators make them prone to island formation which in some devices is used deliberately as exhaust layer between plasma core and the material wall elements around the plasma [15].

Since particle drift extent depends on Larmor radius, electrons stream along the field lines, while ions have larger mass and, hence, larger shift of the drift orbit from the flux surface. This results in an ambipolar field, with the same symmetry as the main magnetic island, to balance the drifts and ensure quasi-neutrality. On the basis of transport particle simulations performed through the guiding center (GC) Hamiltonian code ORBIT [16], a model of electrostatic potential was built up for the island resonating with poloidal/toroidal mode number  $m/n = 0/1$  at the edge of RFX-mod [13], which reproduces the main features of  $E^r$ , such as amplitude and geometry along the toroidal angle  $\varphi$ .

In this Paper we present the first algebraic determination of an ambipolar potential in a tokamak with resonant magnetic perturbation fields applied. We study a circular shaped, high field side limited plasma at the TEXTOR tokamak [17], where a stochastic layer can be generated at the edge, by inducing RMPs through the Dynamic Ergodic Divertor (DED) [18]. The particle transport properties of the magnetic topology in the DED configuration  $m/n = 12/4$  were analyzed using Poincaré plots and by calculating the parallel connection length,  $L_{\parallel}$ , for ions and electrons. The resulting  $L_{\parallel}$  map showed a radial and poloidal modulation, being footprints of the magnetic topology [19]. Results with ORBIT are consistent with maps of connection length made with GOURDON [20]. Previous results have pointed to the formation of island convective cells due to  $E \times B$  flows around magnetic islands [21, 22, 23, 24]. Here, we show for the first time that ambipolar potentials, with the same symmetry as the magnetic island, due to differential drifts of ions and electrons in islands can

account for the radial electric fields responsible for these flows. This potential is the response of the plasma to non-ambipolar fluxes, generated through the breaking of the toroidal symmetry by the 3D fields. The analogy is that of the neoclassical response to magnetic islands in the stellarator [25], where banana trajectories do not close onto themselves on the poloidal plane, due to the 3D distortion [26, 27]. In the case of a stochastic edge, such as in TEXTOR, the tiny details of stochastic layers and fixed points dominate over neoclassical effects in driving transport, and an optimal tool is a guiding-center (GC) code, such as ORBIT [16]. ORBIT has an Hamiltonian formulation of the GC equations of motion, plus the additional capability of describing collisional effects for electrons/ions, through a Monte-Carlo package based on the Boozer-Kuo operator [28]. A heuristic argument for demonstrating the need of a potential  $\Phi$  to balance the asymmetry of electron/ion fluxes in presence of a 3D perturbation, can be derived directly from the GC equations. Specify the equation for electrons and ions [29], and neglect the ripple:

$$\dot{P}_\zeta^{(e)} = \rho_\parallel^e B^2 \partial_\zeta \alpha + \frac{\partial \Phi}{\partial \zeta} \quad (1)$$

$$\dot{P}_\zeta^{(i)} = \rho_\parallel^i B^2 \partial_\zeta \alpha - \frac{\partial \Phi}{\partial \zeta} , \quad (2)$$

where  $\rho_\parallel = mv_\parallel/eB$  is the “parallel” gyro-radius,  $P_\zeta$  is the canonical toroidal momentum, and the magnetic field perturbation is treated as  $\delta \vec{B} = \nabla \times \alpha \vec{B}_0$ . Flux coordinates of Boozer-type  $(\psi_p, \theta, \zeta)$  are used. The meaning of Eqs. (1-2) is that, in presence of a 3D field  $\alpha$ , the toroidal momentum  $P_\zeta$  is no more conserved in time. On the other hand, a larger drift (larger  $\rho_\parallel^i$ ) for ions determines a different response to the symmetry breaking brought about by  $\partial_\zeta \alpha$ , and this different change in  $\dot{P}_\zeta$  must be balanced by the ambipolar potential  $\Phi$ . Subtract (1) from (2)

$$\dot{P}_\zeta^{(i)} - \dot{P}_\zeta^{(e)} = (\rho_\parallel^i - \rho_\parallel^e) B^2 \partial_\zeta \alpha - 2 \frac{\partial \Phi}{\partial \zeta} = 0 , \quad (3)$$

and solve in terms of the potential

$$\frac{\partial \Phi}{\partial \zeta} = \frac{1}{2} (\rho_\parallel^i - \rho_\parallel^e) B^2 \partial_\zeta \alpha . \quad (4)$$

If  $\alpha$  is a single mode,  $\alpha = \alpha_{m,n} \sin(m\theta - n\zeta + \phi)$ , with  $\phi$  phase of the mode, Eq. (4) can be integrated to give

$$\begin{aligned} \Phi(\psi_p, \theta, \zeta) &= \Phi_0(\psi_p) + \frac{1}{2} (\rho_\parallel^i - \rho_\parallel^e) B^2 \alpha_{m,n}(\psi_p) \sin(m\theta - n\zeta + \phi) \\ &= \Phi_0(\psi_p) + \frac{1}{2} (\rho_\parallel^i - \rho_\parallel^e) B^2 \alpha_{m,n}(\psi_p) \sin u , \end{aligned} \quad (5)$$

where  $u$  is the helical angle. This heuristic argument does not catch the overall complexity of the electron and ion motion (full ORBIT simulations are needed), but it shows that, whenever you break the symmetry, this is done differently for electrons and ions, and a balancing  $\Phi$  is needed, which will be modulated as  $\sin u$ . This is independent of the shape of the equilibrium flux surfaces  $\psi_p$ , and it is valid also for slowly rotating islands, where  $u = m\theta - n\zeta + \omega t$ .

The experiments on TEXTOR which are used in this paper have been performed in the L-mode wall limited circular plasmas, with a mode number resonant field

$m/n = 3/1$  produced with the Dynamic Ergodic Divertor (DED). The numerical interpretation was conducted using test particle transport simulations by means of ORBIT. We calculate the particle diffusion coefficients for electrons,  $D_e$ , and ions,  $D_i$ , and develop a model for the ambipolar potential, which describes the two-fluid, plasma response to the RMP. The modeled potential reproduces quite well measurements of plasma potential, performed with a Langmuir sweeping probe, inside of a  $m/n = 4/1$  island which is formed close to the plasma edge. The results show that the development of an electrostatic potential is a general feature of magnetic islands resonating at the plasma edge: moreover, two possible ambipolar solutions are present, which resemble the “ion” and “electron-roots” typical of the  $E^r$  in stellarators [30]. A consequence is that modifying the  $T_e/T_i$  ratio can let the system flip from one solution to the other.

We consider a TEXTOR discharge with static RMP, in the  $m/n = 3/1$  operational mode. In Fig. 1 we show the Poincaré plot of the vacuum magnetic field lines, superimposed to the helical 4/1 flux surfaces,  $\psi_h^{(4,1)}$  [31], used for the computation domain and displayed as blue curves. We can recognize the characteristic magnetic topology of TEXTOR at the edge [20]: in the inner region the last main island chain composed by three conserved structures (green points), in the middle four *remnant islands* (purple points) and in the outermost region the *laminar flux tubes* embedded into the *stochastic fingers*. In Ref. [13] we presented the calculation of the




Figure 1: Poincaré plot of the vacuum magnetic field lines, superimposed to the helical 4/1 flux surfaces,  $\psi_h^{(4,1)}$  (blue curves). The x-axis is the poloidal angle while the y-axis is the normalized radius.

particle diffusion coefficients,  $D$ , in between fixed points, O and X (OP and XP in the remainder of the Letter). Here, we propose again the result as it is complementary to the measurements of plasma potential and the modeled ambipolar potential, that we present below.  $D$  is calculated in an helical domain centered at the  $q = 4$  resonance, ( $r \approx 36$  cm), and bounded by  $\psi_h^{(4,1)}$ , highlighted in orange and light green in Fig. 1, respectively, which can be shifted from the OP towards the XP by varying the phase,  $\phi$ , of  $\psi_h^{(4,1)}$ . We considered temperature  $T_e = 90$  eV and  $T_i = 100$  eV and thermal collisions with a background at density  $n_e = 8.7 \times 10^{12}$  cm $^{-3}$ . These values are chosen to approximate the experiment conditions and calculated through the transport code EMC3-Eirene [32] in unperturbed conditions. The area of the domain is an *Archimedes' serpentine*, namely, a cyclic helical surface generated by the helical motion of a circle, whose area is  $\mathcal{A} = 4\pi^2 b \sqrt{r_s^2 + R^2 q^2}$ . In the formula,  $b$  is the radius of the circle normal to the helix,  $r_s$  is the resonance radius, and  $R$  the major radius.

$D_e$  and  $D_i$  are shown in Fig. 2 as a function of the helical angle  $u_{4,1} = 4\theta - \zeta + \phi$  [7], with  $\theta$  and  $\zeta$  the general poloidal and toroidal angle, respectively ( $\zeta = \varphi - \nu(\psi_p, \theta)$ , with  $\nu$  required to fulfill the straight-field line condition in Boozer coordinates [33]).

Fig. 2 is adapted from Fig. 6 in Ref. [13]. Some  $D_e$  values have been corrected ( $D_e$  curve is smoother), but the overall result does not change.  $D_i$  is rather constant along the path ( $\approx 0.1 \text{ m}^2/\text{s}$ ), while  $D_e$  is larger, with typical values in a stochastic field [34] ( $0.6 \div 40 \text{ m}^2/\text{s}$ ). More important,  $D_e$  is strongly modulated along  $u$  (larger at the XP, lower at the OP), consistently with the  $L_{\parallel}$  simulations in Ref. [19], and the well-known experimental result that the laminar flux tubes (XP of the 4/1 island) are pathways of enhanced electron diffusion [20].




Figure 2:  $D_i$  and  $D_e$  values along the helical flux in between the OP and XP at  $3\pi/2$ . On the x-axis the helical angle  $u = m\theta - n\zeta + \phi$ .

Measurements of plasma potential have been done in the region of the 4/1 island by means of a fast insertable probe located at the low field side of the TEXTOR device [35]. The island is generated as a bifurcation with island opening at a given threshold DED current ( $1.8 \text{ kA}$  for this configuration). Then the island, once generated, is moved poloidally by changing the phase of the DED current from shot to shot and in each shot one radial plunge of the fast reciprocating probe is taken. Fig. 3 shows a colored map of the measured plasma potential  $V_p$  in the  $(r, \theta)$  plane, together with the helical flux surfaces  $\psi_h^{(4,1)}$  and the magnetic field Poincaré map are overplotted to  $V_p$ . A very clear correlation of the  $V_p$  shape with the magnetic topology is found. In particular, the correlation is very strong in the region outside of the last closed flux surface (LCFS), while inside  $V_p$  does not follow exactly the flux surfaces. On the basis of the simulations of  $D$  and the measured  $V_p$  map, we find good reason to assume that the ambipolar potential  $\Phi$  should possess the same geometry as the 4/1 island, similarly to the 0/1 and 1/7 island cases in RFX-mod [7, 13].

We now want to understand the link between the high electron diffusivity and the electric field structure. We will use the algebraic determination of the ambipolar electric field, which is customary in the stellarator community: some examples include FORTEC-3D [36, 37] and EUTERPE-GSRAKE [38], or the linearized drift-kinetic equation, such as in the case of DKES-PENTA [39]. In the chaotic transport calculations with ORBIT it is easier to find a proper analytic form for  $\Phi$  to stick into the GC equations of motion, Eqs (1,2). To do that, we need to mix an experimental radial profile and




Figure 3: Map of the measured plasma potential  $V_p$  as a function of the poloidal angle  $\theta$  and the normalized radius  $r$ . The helical flux surfaces  $\psi_h^{(4,1)}$  (white contours) and the magnetic field Poincaré plot (points) are overlotted.



simulation observations along  $\theta$ , as previously done on RFX-mod [13]. In this way,

$$\Phi(\psi_p, \theta, \zeta) = \Phi_0 \left( f_1 + \frac{1}{2}(f_2 - f_1) \sin(-m\theta + n\zeta + \tilde{\phi}) \right), \quad (6)$$

where

$$f_i(\psi_p) = V_{p,i}^{min} + \frac{1}{2}(V_{p,i}^{max} - V_{p,i}^{min}) \left( 1 - \tanh\left(\frac{\psi_p - \psi_{p,i}}{\Delta\psi_{p,i}}\right) \right), \quad (7)$$

with  $i = (1, 2)$ .  $f_1$  and  $f_2$  are the curves fitting the radial profile of  $V_p$  (normalized to  $\langle V_p \rangle \approx 85$  V in the OP) at the poloidal positions of the XP ( $V_{p,1}^{min} = 0.35$ ,  $V_{p,1}^{max} = 0.94$ ,  $\psi_{p,1} = 0.0145$ ,  $\Delta\psi_{p,1} = 0.0005$ ) and the OP ( $V_{p,2}^{min} = 0.41$ ,  $V_{p,2}^{max} = 1.00$ ,  $\psi_{p,2} = 0.0148$ ,  $\Delta\psi_{p,2} = 0.0003$ ), respectively. By setting  $\Phi_0 = 90$  V (the maximum amplitude in the measurements) and  $\tilde{\phi} = \phi$ , i.e. the same phase of  $\psi_h^{(4,1)}$ , we obtain a model  $\Phi$ , identical to the measured plasma potential, as shown in Fig. 4. This is not surprising, considering the radial modulation of  $\Phi$  which coincides by construction with measurements; but the fact that the poloidal dependence follows the geometry of the island is a striking new result in the tokamak. This behavior was already found instead in the reversed-field pinch RFX-mod [7], and in gyrokinetic simulations in stellarators [38]. In Fig. 5, we map the  $E^r = -\partial\Phi/\partial r$  amplitude together with the flux surfaces  $\psi_h^{(4,1)}$  and magnetic field Poincaré plot, noting that  $E^r$  is modulated *both in the radial and in the poloidal directions*. In particular, a region of large positive  $E^r$ , along the LCFS, can be noticed. This is a confirmation of the well known presence of a positive  $E^r$  in the stochastic edge [40, 41, 42]. But, if we focus on this region, we can note also a modulation in the poloidal angle, strictly linked to the magnetic

Figure 4: Map of the modeled potential,  $\Phi$ , in the  $(\theta, r)$  plane.Figure 5: Map of the modeled  $E^r$  in the  $(\theta, r)$  plane.

topology, too:  $E^r$  has a minimum in between the XP and the OP, and an absolute maximum in correspondence of the XP. On the contrary, right into the OP,  $E^r$  almost vanishes, which is consistent with LHD results [24, 43]. Therefore, the potential well is located near the XP, where the electrons are preferably lost, as shown in Fig. 2 and in Ref. [19]. This rather complicated behavior of  $E^r$  should be accounted for when analyzing data in presence of RMPs [10, 44], since  $E^r$  varies both over  $r$  and  $\theta$ . Our results have been obtained in the case of a static RMP: this solution is valid even in the case of slowly rotating islands, as in the case of RFX [13]. In the case of fast rotating islands (in TEXTOR, with frequency  $\omega > 3$  kHz), it is necessary to account for an inductive correction for  $\Phi$ , due to the term  $\partial_t \alpha \vec{B}$  that adds to the expression of Eq. (5). The complete expression for this inductive correction can be found in [45], but it results in a simple Doppler shift of the potential  $\Phi$  in the rotating frame of the island, which is consistent with the results by Stoschus *et al.* [6].

As a final test, we check the ambipolarity of  $\Phi$  by keeping  $\tilde{\phi} = \phi$  (maximum potential at the OP) and evaluating the electron and ion fluxes as a function of




Figure 6: Ion (blue) and electron (red) fluxes as a function of  $\Phi_0$  and the maximum  $E^r$ .

the maximum potential  $\Phi_0$ : this is the algebraic way of determining the ambipolar solution, used in the stellarator community [30]. Even if the method is well-known since a long time, and its possible extension to a symmetry-breaking perturbation has already been mentioned elsewhere [25], this is the first time that the calculation has been fully carried on. To do this, we adapted ORBIT guiding center equations [16] to correctly express electron drifts. We evaluate fluxes at  $\psi_h$ , with source at  $q = 4$ . In Fig. 6 we plot the ion and electron fluxes as a function of  $\Phi_0$  and the maximum  $E^r$ . The two curves show two roots, similarly to stellarators [30]: an unstable ion-root at  $E^r < 0$  ( $\sim -150$  V) and a stable electron-root at  $E^r > 0$  ( $\sim 120$  V), where the latter is found for a positive potential consistently with the experimental findings (see Fig. 3). This shows that two solutions are possible: one with the potential well (=maximum  $E^r$ ) at the XP of the RMP (stable “electron root”, which is the solution found in experiment), and the other with the potential well at the OP (unstable, “ion root”). Here the name for the roots follows the usual meaning given in stellarators, where “electron root” means the ambipolar root for the fastest particle involved in radial transport. With the  $T_e/T_i \sim 0.9$  ratio of TEXTOR, the electron root is favored, but in principle it is possible, by acting on the  $T_e/T_i$  ratio, to make the system flip to the ion root. A sensitivity scan on this point can be done with ORBIT, by varying  $T_e/T_i$  (increasing  $T_i$ ), which would correspond to applying ion cyclotron resonance heating (ICRH) in an experiment. We show the  $D_e/D_i$  ratio as a function of  $T_e/T_i$  in Fig. 7, with the diffusion coefficients calculated in the OP and XP.  $D_e/D_i$  decreases by increasing  $T_i$  similarly in the OP and XP, moving from the TEXTOR experimental condition, marked as a vertical red line in the picture. In the OP the system flips to the ion root ( $D_e/D_i < 1$ ) for  $T_e/T_i \lesssim 0.5$ . The opposite is seen experimentally in stellarators, where the electron root can be induced actively by electron cyclotron resonance heating (ECRH) [30]. Indeed, experimental results in the ASDEX-U and FTU tokamaks show that disruptions can be mitigated by ECRH targeted on the 2/1 island [46]. We speculate that ECRH can modify the  $E^r$  distribution in the edge, and in this way the overall magnetohydrodynamics stability at the edge. In principle, this can be also a way of overcoming the density limit, which critically depends on the  $E^r$  pattern, at least in the RFP [12]. Finally, it should be worth doing experiments






Figure 7:  $D_e/D_i$  as a function of  $T_e/T_i$  calculated in the OP and XP.

of ECRH in conjunction with RMP, to assess the role of  $E^r$  on plasma stability with respect to the so-called edge localized modes [14].

In summary, we analyzed the local radial particle transport along a helical path from the OP through the XP of an  $m/n = 4/1$  remnant island, created near the edge of TEXTOR. Electron diffusion is strongly modulated (larger at the XP, lower at the OP), which requires a large electrostatic potential to ensure quasi-neutrality. We developed a 3D model for the ambipolar potential on the basis of the geometry of the remnant island: the resulting  $E^r$  shows a large positive value near the LCFS, confirming a well known result in the RMP tokamak community. The mechanism of ambipolarity shows two possible solutions ("roots"), which suggests a way of acting on the edge  $E^r$  through additional heating.

This project has received funding from the European Union Horizon 2020 research and innovation programme under grant agreement number 633053, and in part from the U.S. Department of Energy Grant DE-AC02-09CH11466.

## References

- [1] Frolov V P 2012 *Phys. Rev. D* **85**(2) 024020 URL <http://link.aps.org/doi/10.1103/PhysRevD.85.024020>
- [2] Meier D L 2003 *New Astronomy Reviews* **47** 667 – 672 ISSN 1387-6473 URL <http://www.sciencedirect.com/science/article/pii/S1387647303001209>
- [3] Evans T E, Moyer R A, Burrell K H, Fenstermacher M E, Joseph I, Leonard A W, Osborne T H, Porter G D, Schaffer M J, Snyder P B, Thomas P R, Watkins J G and West W P 2006 *Nature Phys.* **2** 419–423 URL <http://dx.doi.org/10.1038/nphys312>
- [4] Schmitz O, Evans T E, Fenstermacher M E, Unterberg E A, Austin M E, Bray B D, Brooks N H, Frerichs H, Groth M, Jakubowski M W, Lasnier C J, Lehnen M, Leonard A W, Mordijck S, Moyer R A, Osborne T H, Reiter D, Samm U, Schaffer M J, Unterberg B and West W P (the DIII-D and TEXTOR Research Teams) 2009 *Phys. Rev. Lett.* **103** 165005 URL <http://link.aps.org/doi/10.1103/PhysRevLett.103.165005>
- [5] Moyer R, Zeeland M V, Orlov D, Wingen A, Evans T, Ferraro N, Hanson J, Nazikian R, Wade M and Zeng L 2012 *Nuclear Fusion* **52** 123019 URL <http://stacks.iop.org/0029-5515/52/i=12/a=123019>
- [6] Stoschus H, Schmitz O, Frerichs H, Reiser D, Jakubowski M, Unterberg B, Lehnen M, Reiter D, Samm U and the TEXTOR team 2012 *Nuclear Fusion* **52** 083002 URL <http://stacks.iop.org/0029-5515/52/i=8/a=083002>
- [7] Vianello N, Spizzo G, Agostini M, Scarin P, Carraro L, Cavazzana R, De Masi G, Martines E, Momo B, Rea C, Spagnolo S, Spolaore M, Zuin M and the RFX-Mod Team 2013 *Nuclear Fusion* **53** 073025 URL <http://stacks.iop.org/0029-5515/53/i=7/a=073025>

- [8] Feng Y, Kobayashi M, Lunt T and Reiter D 2011 *Plasma Physics and Controlled Fusion* **53** 024009 URL <http://stacks.iop.org/0741-3335/53/i=2/a=024009>
- [9] Kamiya K, Ida K, Yoshinuma M, Suzuki C, Suzuki Y, Yokoyama M and the LHD Experiment Group 2013 *Nuclear Fusion* **53** 013003 URL <http://stacks.iop.org/0029-5515/53/i=1/a=013003>
- [10] Mordijck S, Moyer R, Ferraro N, Wade M and Osborne T 2014 *Nuclear Fusion* **54** 082003 URL <http://stacks.iop.org/0029-5515/54/i=8/a=082003>
- [11] Scarin P, Vianello N, Agostini M, Spizzo G, Spolaore M, Zuin M, Cappello S, Carraro L, Cavazzana R, De Masi G, Martinez E, Moresco M, Munaretto S, Puiatti M E, Valisa M and the RFX-mod Team 2011 *Nuclear Fusion* **51** 073002 URL <http://stacks.iop.org/0029-5515/51/i=7/a=073002>
- [12] Puiatti M, Valisa M, Agostini M, Auriemma F, Bonomo F, Carraro L, Fassina A, Gobbin M, Lorenzini R, Momo B, Scaggion A, Zaniol B, Alfieri A, Apolloni L, Baruzzo M, Bolzonella T, Bonfiglio D, Canton A, Cappello S, Cavazzana R, Bello S D, Masi G D, Escande D, Franz P, Gazza E, Guo S, Innocente P, Marchiori G, Marrelli L, Martin P, Martines E, Martini S, Menmuir S, Novello L, Paccagnella R, Piovesan P, Piron L, Predebon I, Ruzzon A, Sattin F, Scarin P, Soppelsa A, G Spizzo, Spagnolo S, Spolaore M, Terranova D, Veranda M, Vianello N, Zanca P, Zanutto L and Zuin M 2011 *Nuclear Fusion* **51** 073038 URL <http://stacks.iop.org/0029-5515/51/i=7/a=073038>
- [13] Spizzo G, Vianello N, White R B, Abdullaev S S, Agostini M, Cavazzana R, Ciaccio G, Puiatti M E, Scarin P, Schmitz O, Spolaore M, Terranova D and RFX and TEXTOR Teams 2014 *Physics of Plasmas (1994-present)* **21** 056102 URL <http://scitation.aip.org/content/aip/journal/pop/21/5/10.1063/1.4872173>
- [14] Lang P, Loarte A, Saibene G, Baylor L, Becoulet M, Cavinato M, Clement-Lorenzo S, Daly E, Evans T, Fenstermacher M, Gribov Y, Horton L, Lowry C, Martin Y, Neubauer O, Oyama N, Schaffer M, Stork D, Suttrop W, Thomas P, Tran M, Wilson H, Kavin A and Schmitz O 2013 *Nuclear Fusion* **53** 043004 URL <http://stacks.iop.org/0029-5515/53/i=4/a=043004>
- [15] Renner H, Boscary J, Erckmann V, Greuner H, Grote H, Sapper J, Speth E, Wesner F, Wanner M and Team W X 2000 *Nuclear Fusion* **40** 1083 URL <http://stacks.iop.org/0029-5515/40/i=6/a=306>
- [16] White R B and Chance M S 1984 *Physics of Fluids* **27** 2455–2467 URL <http://link.aip.org/link/?PFL/27/2455/1>
- [17] Neubauer O, Czymek G, Giesen B, Hüttemann P W, Sauer M, Schalt W and Schruoff J 2005 *Fus. Sci. Technol.* **47** 76–86 URL <http://epubs.ans.org/?a=689>
- [18] Abdullaev S 2014 *Magnetic Stochasticity in Magnetically Confined Fusion Plasmas (Springer Series on Atomic, Optical, and Plasma Physics vol 78)* (Springer Cham Heidelberg New York Dordrecht London) ISBN 978-3-319-01889-8 URL <http://www.springer.com/physics/atomic,+molecular,+optical+%26+plasma+physics/book/978-3-319-01889-8>
- [19] Ciaccio G, Schmitz O, Abdullaev S, Frerichs H, Agostini M, Scarin P, Spizzo G, Vianello N and White R B 2014 *Nuclear Fusion* **54** 064008 URL <http://stacks.iop.org/0029-5515/54/i=6/a=064008>
- [20] Schmitz O, Jakubowski M, Frerichs H, Harting D, Lehnen M, Unterberg B, Abdullaev S, Brezinsek S, Classen I, Evans T, Feng Y, Finken K, Kantor M, Reiter D, Samm U, Schweer B, Sergienko G, Spakman G, Tokar M, Uzel E, Wolf R and the TEXTOR Team 2008 *Nuclear Fusion* **48** 024009 URL <http://stacks.iop.org/0029-5515/48/i=2/a=024009>
- [21] Evans T E *et al.* 1987 *Proc. of the 14<sup>th</sup> EPS Conference on Plasma Physics* vol 11D ed Engelmann F and Alvarez Rivas J L (Madrid, Spain: European Physical Society (Petit Lancy)) pp 770–773
- [22] McCool S, Wootton A, Kotschenreuther M, Audemir A, Bravenec R, DeGrassie J, Evans T, Hickok R, Richards B, Rowan W and Schoch P 1990 *Nuclear Fusion* **30** 167 URL <http://stacks.iop.org/0029-5515/30/i=1/a=014>
- [23] Takamura S, Ohnishi N, Yamada H and Okuda T 1987 *Physics of Fluids (1958-1988)* **30** 144–147 URL <http://scitation.aip.org/content/aip/journal/pof1/30/1/10.1063/1.866180>
- [24] Ida K and *et al.* 2004 *Nuclear Fusion* **44** 290 URL <http://stacks.iop.org/0029-5515/44/i=2/a=010>
- [25] Callen J 2011 *Nuclear Fusion* **51** 094026 URL <http://stacks.iop.org/0029-5515/51/i=9/a=094026>
- [26] Shaing K C 2002 *Physics of Plasmas (1994-present)* **9** 3470–3475 URL <http://scitation.aip.org/content/aip/journal/pop/9/8/10.1063/1.1491533>
- [27] Shaing K, Hegna C, Callen J and Houlberg W 2003 *Nuclear Fusion* **43** 258 URL <http://stacks.iop.org/0029-5515/43/i=4/a=306>
- [28] Boozer A H 1981 *Physics of Fluids* **24** 1999–2003 URL

- <http://link.aip.org/link/?PFL/24/1999/1>
- [29] White R B, Spizzo G and Gobbin M 2013 *Plasma Physics and Controlled Fusion* **55** 115002 URL <http://stacks.iop.org/0741-3335/55/i=11/a=115002>
  - [30] Hastings D, Houlberg W and Shaing K 1985 *Nuclear Fusion* **25** 445 URL <http://stacks.iop.org/0029-5515/25/i=4/a=005>
  - [31] Ciaccio G, Veranda M, Bonfiglio D, Cappello S, Spizzo G, Chacon L and White R B 2013 *Physics of Plasmas* **20** 062505 (pages 12) URL <http://link.aip.org/link/?PHP/20/062505/1>
  - [32] Kobayashi M, Feng Y, Sardei F, Reiter D, Finken K and Reiser D 2004 *Nuclear Fusion* **44** S64 URL <http://stacks.iop.org/0029-5515/44/i=6/a=S07>
  - [33] White R B 2006 *The theory of toroidally confined plasmas* (57 Shelton Street, Covent Garden, London WC2H 9HE: Imperial College Press) p 42 2nd ed ISBN 1860946399 URL <http://www.icpress.co.uk/physics/p440.html>
  - [34] Spizzo G, White R B, Cappello S and Marrelli L 2009 *Plasma Physics and Controlled Fusion* **51** 124026 URL <http://stacks.iop.org/0741-3335/51/i=12/a=124026>
  - [35] Xu Y, Schoor M V, Weynants R, Jachmich S, Vergote M, Jakubowski M, Beyer P, Mitri M, Schweer B, Reiser D, Unterberg B, Finken K, Lehnen M, Jaspers R and the TEXTOR Team 2007 *Nuclear Fusion* **47** 1696 URL <http://stacks.iop.org/0029-5515/47/i=12/a=008>
  - [36] Satake S, Idomura Y, Sugama H and Watanabe T H 2010 *Computer Physics Communications* **181** 1069–1076 ISSN 0010-4655 URL <http://www.sciencedirect.com/science/article/pii/S0010465510000536>
  - [37] Matsuoka S, Satake S, Yokoyama M, Wakasa A and Murakami S 2011 *Physics of Plasmas* **18** 032511 (pages 11) URL <http://link.aip.org/link/?PHP/18/032511/1>
  - [38] García-Regaña J M, Kleiber R, Beidler C D, Turkin Y, Maaßberg H and Helander P 2013 *Plasma Physics and Controlled Fusion* **55** 074008 URL <http://stacks.iop.org/0741-3335/55/i=7/a=074008>
  - [39] Lore J, Guttentfelder W, Briesemeister A, Anderson D T, Anderson F S B, Deng C B, Likin K M, Spong D A, Talmadge J N and Zhai K 2010 *Physics of Plasmas (1994-present)* **17** 056101 URL <http://scitation.aip.org/content/aip/journal/pop/17/5/10.1063/1.3300465>
  - [40] Hess W R, DeMichelis C, Mattioli M, Clairet F, Druetta M, Grosman A, Guirlet R, Hutter T, Lasalle J and Monier-Garbet P 1995 *Plasma Physics and Controlled Fusion* **37** 951 URL <http://stacks.iop.org/0741-3335/37/i=9/a=003>
  - [41] Rozhansky V, Kaveeva E, Molchanov P, Veselova I, Voskoboinikov S, Coster D, Kirk A, Liso S and Nardon E 2010 *Nuclear Fusion* **50** 034005 URL <http://stacks.iop.org/0029-5515/50/i=3/a=034005>
  - [42] Xu Y, Weynants R R, Jachmich S, Van Schoor M, Vergote M, Peleman P, Jakubowski M W, Mitri M, Reiser D, Unterberg B and Finken K H (the TEXTOR team) 2006 *Phys. Rev. Lett.* **97**(16) 165003 URL <http://link.aps.org/doi/10.1103/PhysRevLett.97.165003>
  - [43] Ida K, Ohyabu N, Morisaki T, Nagayama Y, Inagaki S, Itoh K, Liang Y, Narihara K, Kostrioukov A Y, Peterson B J, Tanaka K, Tokuzawa T, Kawahata K, Suzuki H and Komori A (LHD Experimental Group) 2001 *Phys. Rev. Lett.* **88**(1) 015002 URL <http://link.aps.org/doi/10.1103/PhysRevLett.88.015002>
  - [44] Coenen J, Schmitz O, Unterberg B, Clever M, Jakubowski M, Samm U, Schweer B, Stoschus H, Tokar M and the TEXTOR-Team 2011 *Nuclear Fusion* **51** 063030 URL <http://stacks.iop.org/0029-5515/51/i=6/a=063030>
  - [45] White R B 2006 *The theory of toroidally confined plasmas* (57 Shelton Street, Covent Garden, London WC2H 9HE: Imperial College Press) p 242 2nd ed ISBN 1860946399 URL <http://www.icpress.co.uk/physics/p440.html>
  - [46] Esposito B, Granucci G, Smeulders P, Nowak S, Martín-Solís J R and Gabel-lerri L (FTU and ECRH teams) 2008 *Phys. Rev. Lett.* **100**(4) 045006 URL <http://link.aps.org/doi/10.1103/PhysRevLett.100.045006>